



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Vernon Bailey. It has already been recorded incidentally (Cooke, Bull. U. S. Dept. Agric., No. 292, October 25, 1915, p. 47), but owing to its importance it seems worthy of special notice in a place more accessible to ornithologists generally. It represents the easternmost record of *Larus canus brachyrhynchus*, and the only really interior occurrence of the species in the United States. For the change of the name of this bird from *Larus brachyrhynchus* to *Larus canus brachyrhynchus* see 'The Auk,' XXXVI, No. 1, January, 1919, p. 83.—HARRY C. OBERHOLSER, Washington, D. C.

***Polysticta Eyton* versus *Stelleria Bonaparte*.**—Mr. G. M. Mathews has recently (Austral Avian Record, III, No. 5, December 28, 1917, p. 123) advocated the use of the generic name *Stelleria* Bonaparte for the species now known as *Polysticta stelleri* (Pallas). The term *Stelleria* is, of course, as he shows, not debarred from employment in zoology by the previous use of *Stelleria* in botany; but he has apparently overlooked the fact that *Polysticta* is not preoccupied, since *Polysticta* Smith (Illust. South Afr. Zoöl.), June [or later], 1836, does not invalidate *Polysticta Eyton* (Catal. Brit. Birds), April, 1836, a fact to which Dr. C. W. Richmond long ago (Proc. Biol. Soc. Wash. XVI, September 30, 1903, p. 128) called attention. It is evident, therefore, that the name of Steller's Eider should remain *Polysticta stelleri* (Pallas).—HARRY C. OBERHOLSER, Washington, D. C.

**Further Record of the European Widgeon at Madison, Wis.**—On April 14, 1918, in the wide-water at the head of Lake Waubesa, four miles south of Madison, I was able to identify unmistakably a typical specimen of the European Widgeon (*Mareca penelope*) that was in the company of seventeen Baldpates (*Mareca americana*). The bird was drawn so close by my 40-power telescope that it covered one-third of the field and allowed close study.

It may be of further interest to restate the substance of a note submitted by Mr. A. W. Schorger to the January, 1918 issue of 'The Auk' in regard to the recent appearance of the European Widgeon in the vicinity of Madison. On April 22, 1917, a specimen was discovered by Mr. Schorger on the Hammersley Marsh in company with about thirty Baldpates and a few other ducks. It remained at least four days and was seen by me at close range on three occasions, the last being on the 26th. On the 28th Mr. George H. Jenkins observed a specimen, perhaps the same, among a flock of Baldpates on the Yahara Marshes ten miles distant.—WARNER TAYLOR, Madison, Wisconsin.

**A Late Record for *Rallus elegans* for Maine.**—November 22, 1909, Mr. A. G. Dorr, Bucksport, Me., collected and sent me in the flesh a fine male specimen of the King Rail. It measured as follows: length, 16.30; wing, 6.75; tail, 2.10; tarsus, 2.34 and bill, 2.40 inches. It was marked above with brownish-black and olive-brown feather edging; light throat and rufous-cinnamon, breast and flanks fuscous, distinctly barred with